

Linear Odd Poisson Bracket on Grassmann Algebra

Vyacheslav A. Soroka*

*Kharkov Institute of Physics and Technology
310108 Kharkov, Ukraine*

Abstract

A linear odd Poisson bracket realized solely in terms of Grassmann variables is suggested. It is revealed that with the bracket, corresponding to a semi-simple Lie group, both a Grassmann-odd Casimir function and invariant (with respect to this group) nilpotent differential operators of the first, second and third orders are naturally related and enter into a finite-dimensional Lie superalgebra.

1. Recently a linear degenerate odd Poisson bracket built only of Grassmann variables has been introduced [1]. It was constructed for this bracket, in contrast with the non-degenerate odd bracket having the only Grassmann-odd nilpotent differential Δ -operator of the second order, at once three Grassmann-odd nilpotent Δ -like differential operators of the first, the second and the third orders with respect to Grassmann derivatives. It was also shown that these Δ -like operators together with a Grassmann-odd nilpotent Casimir function of this degenerate odd bracket form a finite-dimensional Lie superalgebra. Following to [2], in the present report we extend the above-mentioned results to the case of an arbitrary linear odd Poisson bracket, which is also realized solely in terms of the Grassmann variables and corresponds to a semi-simple Lie group.

2. There is a well-known linear even Poisson bracket given in terms of the commuting (Grassmann-even) variables X_α ($g(X_\alpha) = 0$)

$$\{X_\alpha, X_\beta\}_0 = c_{\alpha\beta}^\gamma X_\gamma, \quad (\alpha, \beta, \gamma = 1, \dots, N), \quad (1)$$

where $c_{\alpha\beta}^\gamma$ are Grassmann-even ($g(c_{\alpha\beta}^\gamma) = 0$) constants which, because of the main properties of the even Poisson bracket:

$$\{A, B + C\}_0 = \{A, B\}_0 + \{A, C\}_0, \quad (2)$$

$$g(\{A, B\}_0) = g(A) + g(B) \pmod{2}, \quad (3)$$

*E-mail: vsoroka@kipt.kharkov.ua

$$\{A, B\}_0 = -(-1)^{g(A)g(B)} \{B, A\}_0 , \quad (4)$$

$$\sum_{(ABC)} (-1)^{g(A)g(C)} \{A, \{B, C\}_0\}_0 = 0 , \quad (5)$$

$$\{A, BC\}_0 = \{A, B\}_0 C + (-1)^{g(A)g(B)} B \{A, C\}_0 , \quad (6)$$

are antisymmetric in the two lower indices

$$c_{\alpha\beta}{}^\gamma = -c_{\beta\alpha}{}^\gamma \quad (7)$$

and obey the conditions

$$c_{\alpha\lambda}{}^\delta c_{\beta\gamma}{}^\lambda + c_{\beta\lambda}{}^\delta c_{\gamma\alpha}{}^\lambda + c_{\gamma\lambda}{}^\delta c_{\alpha\beta}{}^\lambda = 0 . \quad (8)$$

A sum with the symbol (ABC) in (5) means a summation over cyclic permutations of the quantities A, B, C . In relations (2)-(6) A, B, C are functions of the variables X_α and $g(A)$ is a Grassmann parity of the quantity A . The linear even bracket (1) plays a very important role in the theory of Lie groups, Lie algebras, their representations and applications (see, for example, [3, 4]). The bracket (1) can be realized in a canonical even Poisson bracket

$$\{A, B\}_0 = A \sum_{\alpha=1}^N \left(\overleftarrow{\partial}_{q^\alpha} \overrightarrow{\partial}_{p_\alpha} - \overleftarrow{\partial}_{p_\alpha} \overrightarrow{\partial}_{q^\alpha} \right) B ,$$

where $\overleftarrow{\partial}$ and $\overrightarrow{\partial}$ are the right and left derivatives and $\partial_{x^A} \equiv \frac{\partial}{\partial x^A}$, on the following bilinear functions of coordinates q^α and momenta p_α

$$X_\alpha = c_{\alpha\beta}{}^\gamma q^\beta p_\gamma$$

if $c_{\alpha\beta}{}^\gamma$ satisfy the conditions (7), (8) for the structure constants of a Lie group.

As in the Lie algebra case, we can define a symmetric Cartan-Killing tensor

$$g_{\alpha\beta} = g_{\beta\alpha} = c_{\alpha\gamma}{}^\lambda c_{\beta\lambda}{}^\gamma \quad (9)$$

and verify with the use of relations (8) an anti-symmetry property of a tensor

$$c_{\alpha\beta\gamma} = c_{\alpha\beta}{}^\delta g_{\delta\gamma} = -c_{\alpha\gamma\beta} . \quad (10)$$

By assuming that the Cartan-Killing metric tensor is non-degenerate $\det(g_{\alpha\beta}) \neq 0$ (this case corresponds to the semi-simple Lie group), we can define an inverse tensor $g^{\alpha\beta}$

$$g^{\alpha\beta} g_{\beta\gamma} = \delta_\gamma^\alpha , \quad (11)$$

with the help of which we are able to build a quantity

$$C = X_\alpha X_\beta g^{\alpha\beta} ,$$

that, in consequence of relation (10), is for the bracket (1) a Casimir function which annihilates the bracket (1) and is an invariant of the Lie group with the structure constants $c_{\alpha\beta}{}^\gamma$ and the generators T_α

$$\{X_\alpha, C\}_0 = c_{\alpha\beta}{}^\gamma X_\gamma \partial_{X_\beta} C = T_\alpha C = 0 .$$

3. Now let us replace in expression (1) the commuting variables X_α by Grassmann variables Θ_α ($g(\Theta_\alpha) = 1$). Then we obtain a binary composition

$$\{\Theta_\alpha, \Theta_\beta\}_1 = c_{\alpha\beta}{}^\gamma \Theta_\gamma , \quad (12)$$

which, due to relations (7) and (8), meets all the properties of the odd Poisson brackets:

$$\{A, B + C\}_1 = \{A, B\}_1 + \{A, C\}_1 , \quad (13)$$

$$g(\{A, B\}_1) = g(A) + g(B) + 1 \pmod{2} , \quad (14)$$

$$\{A, B\}_1 = -(-1)^{(g(A)+1)(g(B)+1)} \{B, A\}_1 , \quad (15)$$

$$\sum_{(ABC)} (-1)^{(g(A)+1)(g(C)+1)} \{A, \{B, C\}_1\}_1 = 0 , \quad (16)$$

$$\{A, BC\}_1 = \{A, B\}_1 C + (-1)^{(g(A)+1)g(B)} B \{A, C\}_1 . \quad (17)$$

It is surprising enough that the odd bracket can be defined solely in terms of the Grassmann variables as well as an even Martin bracket [5]. On the following bilinear functions of canonical variables commuting q^α and Grassmann θ_α

$$\Theta_\alpha = c_{\alpha\beta}{}^\gamma q^\beta \theta_\gamma$$

a canonical odd Poisson bracket

$$\{A, B\}_1 = A \sum_{\alpha=1}^N \left(\overleftarrow{\partial}_{q^\alpha} \overrightarrow{\partial}_{\theta_\alpha} - \overleftarrow{\partial}_{\theta_\alpha} \overrightarrow{\partial}_{q^\alpha} \right) B$$

is reduced to the bracket (12) providing that $c_{\alpha\beta}{}^\gamma$ obey the conditions (7), (8).

On functions A, B of Grassmann variables Θ_α the bracket (12) has the form

$$\{A, B\}_1 = A \overleftarrow{\partial}_{\Theta_\alpha} c_{\alpha\beta}{}^\gamma \Theta_\gamma \overrightarrow{\partial}_{\Theta_\beta} B ,$$

The bracket (12) can be either degenerate or non-degenerate in the dependence on whether the matrix $c_{\alpha\beta}{}^\gamma \Theta_\gamma$ in the indices α, β is degenerate or not. Raising and lowering of the indices α, β , the non-degenerate metric tensors (9), (11) relate with each other the adjoint and co-adjoint representations which are equivalent for a semi-simple Lie group

$$\Theta^\alpha = g^{\alpha\beta} \Theta_\beta , \quad \partial_{\Theta^\alpha} = g_{\alpha\beta} \partial_{\Theta_\beta} .$$

Hereafter only the non-degenerate metric tensors (11) will be considered.

4. By contracting the indices in a product of the Grassmann variables with the upper indices and of the successive Grassmann derivatives, respectively, with the lower indices in (8), we obtain the relations

$$\Theta^\alpha \Theta^\beta (c_{\alpha\beta}^\lambda c_{\lambda\gamma}^\delta + 2c_{\gamma\alpha}^\lambda c_{\lambda\beta}^\delta) = 0, \quad \Theta^\alpha \Theta^\beta \Theta^\gamma c_{\alpha\beta}^\lambda c_{\lambda\gamma}^\delta = 0, \quad (18a, b)$$

$$(c_{\alpha\beta}^\lambda c_{\lambda\gamma}^\delta + 2c_{\gamma\alpha}^\lambda c_{\lambda\beta}^\delta) \partial_{\Theta_\alpha} \partial_{\Theta_\beta} = 0, \quad c_{\alpha\beta}^\lambda c_{\lambda\gamma}^\delta \partial_{\Theta_\alpha} \partial_{\Theta_\beta} \partial_{\Theta_\gamma} = 0, \quad (19a, b)$$

which will be used later on many times. In particular, taking into account relation (18b), we can verify that the linear odd bracket (12) has the following Grassmann-odd nilpotent Casimir function

$$\Delta_{+3} = \frac{1}{\sqrt{3!}} \Theta^\alpha \Theta^\beta \Theta^\gamma c_{\alpha\beta\gamma}, \quad (\Delta_{+3})^2 = 0, \quad (20)$$

which is an invariant of the Lie group

$$\{\Theta_\alpha, \Delta_{+3}\}_1 = \Theta_\gamma c_{\alpha\beta}^\gamma \partial_{\Theta_\beta} \Delta_{+3} = S_\alpha \Delta_{+3} = 0 \quad (21)$$

with the generators S_α obeying the Lie algebra permutation relations¹

$$[S_\alpha, S_\beta] = c_{\alpha\beta}^\gamma S_\gamma. \quad (22)$$

It is a well-known fact that, in contrast with the even Poisson bracket, the non-degenerate odd Poisson bracket has one Grassmann-odd nilpotent differential Δ -operator of the second order, in terms of which the main equation has been formulated in the Batalin-Vilkovisky scheme [6, 7, 8, 9, 10, 11] for the quantization of gauge theories in the Lagrangian approach. In a formulation of Hamiltonian dynamics by means of the odd Poisson bracket with the help of a Grassmann-odd Hamiltonian $\bar{H}$ ($g(\bar{H}) = 1$) [12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23] this Δ -operator plays also a very important role being used to distinguish the Hamiltonian dynamical systems, for which the Liouville theorem is valid $\Delta \bar{H} = 0$, from those ones, for which this theorem takes no place $\Delta \bar{H} \neq 0$ [1]².

Now let us try to build the Δ -operator for the linear odd bracket (12). It is remarkable that, in contrast with the canonical odd Poisson bracket having the only Δ -operator of the second order, we are able to construct at once three Δ -like Grassmann-odd nilpotent operators which are differential operators of the first, the second and the third orders respectively

$$\Delta_{+1} = \frac{1}{\sqrt{2}} \Theta^\alpha \Theta^\beta c_{\alpha\beta}^\gamma \partial_{\Theta_\gamma}, \quad (\Delta_{+1})^2 = 0; \quad (23)$$

¹Note that below $[A, B] = AB - BA$ and $\{A, B\} = AB + BA$.

²Note also applications of the odd bracket to the integrability problem [24, 25, 26]

$$\Delta_{-1} = \frac{1}{\sqrt{2}} \Theta_\gamma c_{\alpha\beta}{}^\gamma \partial_{\Theta_\alpha} \partial_{\Theta_\beta} , \quad (\Delta_{-1})^2 = 0 ; \quad (24)$$

$$\Delta_{-3} = \frac{1}{\sqrt{3!}} c_{\alpha\beta\gamma} \partial_{\Theta_\alpha} \partial_{\Theta_\beta} \partial_{\Theta_\gamma} , \quad (\Delta_{-3})^2 = 0 . \quad (25)$$

The nilpotency of the operators Δ_{+1} and Δ_{-1} is a consequence of relations (18b) and (19b). The operator Δ_{+1} is proportional to the second term in a BRST charge

$$Q = \Theta^\alpha G_\alpha - \frac{1}{2} \Theta^\alpha \Theta^\beta c_{\alpha\beta}{}^\gamma \partial_{\Theta^\gamma} ,$$

where Θ^α and ∂_{Θ^α} represent the operators for ghosts and antighosts respectively. Q itself will be proportional to the operator Δ_{+1} if we take the representation S_α (21) for group generators G_α . The operator Δ_{-1} , related with the divergence of a vector field $\{\Theta_\alpha, A\}_1$

$$\partial_{\Theta_\alpha} \{\Theta_\alpha, A\}_1 = \partial_{\Theta_\alpha} S_\alpha A = -\sqrt{2} \Delta_{-1} A ,$$

is proportional to the true Δ -operator for the bracket (12).

It is also interesting to reveal that these Δ -like operators together with the Casimir function Δ_{+3} (20) are closed into the finite-dimensional Lie superalgebra, in which the anticommuting relations between the quantities Δ_λ ($\lambda = -3, -1, +1, +3$) (20), (23)-(25) with the nonzero right-hand side are

$$\{\Delta_{-1}, \Delta_{+1}\} = Z , \quad (26)$$

$$\{\Delta_{-3}, \Delta_{+3}\} = N - 3Z , \quad (27)$$

where

$$N = -c^{\alpha\beta\gamma} c_{\alpha\beta\gamma}$$

is a number of values for the indices α, β, γ ($\alpha, \beta, \gamma = 1, \dots, N$) and

$$Z = D - K \quad (28)$$

is a central element of this superalgebra

$$[Z, \Delta_\lambda] = 0 , \quad (\lambda = -3, -1, +1, +3) . \quad (29)$$

In (28)

$$D = \Theta^\alpha \partial_{\Theta^\alpha} \quad (30)$$

is a "dilatation" operator for the Grassmann variables Θ_α , which distinguishes the Δ_λ -operators with respect to their uniformity degrees in Θ

$$[D, \Delta_\lambda] = \lambda \Delta_\lambda , \quad (\lambda = -3, -1, +1, +3) \quad (31)$$

and is in fact a representation for a ghost number operator, and the quantity K has the form

$$K = \frac{1}{2} \Theta^\alpha \Theta^\beta c_{\alpha\beta}^\lambda c_{\lambda\gamma\delta} \partial_{\Theta_\gamma} \partial_{\Theta_\delta} . \quad (32)$$

The operator Z is also a central element of the Lie superalgebra which contains both the operators Δ_λ (20), (23)-(25), Z (28) and the operator D (30)

$$[Z, D] = 0 . \quad (33)$$

We can add to this superalgebra the generators S_α (21) with the following commutation relations:

$$[S_\alpha, \Delta_\lambda] = 0 , \quad [S_\alpha, Z] = 0 , \quad [S_\alpha, D] = 0 , \quad (34)$$

which indicate that both the Casimir function Δ_{+3} and the operators Δ_λ ($\lambda = -3, -1, +1$), Z and D are invariants of the Lie group with the generators S_α . In order to prove the permutation relations for the Lie superalgebra (20)-(34), we have to use relations (18) and (19). Note that the central element Z (28) coincides with the expression for a quadratic Casimir operator of the Lie algebra (22) for the generators S_α given in the representation (21)

$$S_\alpha S_\beta g^{\alpha\beta} = Z . \quad (35)$$

5. Thus, we see that both the even and odd linear Poisson brackets are internally inherent in the Lie group with the structure constants subjected to conditions (7) and (8). However, only for the linear odd Poisson bracket realized in terms of the Grassmann variables and only in the case when this bracket corresponds to the semi-simple Lie group, there exists the Lie superalgebra (20)-(34) for the Δ -like operators of this bracket.

Note that in the case of the degenerate Cartan-Killing metric tensor (9), relation (10) remains valid and we can construct only two Δ -like Grassmann-odd nilpotent operators: Δ_{-1} (24) and Δ_{-3} (25), which satisfy the trivial anticommuting relation

$$\{\Delta_{-1}, \Delta_{-3}\} = 0 .$$

Note also that anticommuting relations for the operators

$$\overset{i}{\Delta}_{-1} = \frac{1}{\sqrt{2}} \Theta_\gamma c_{\alpha\beta}^i{}^\gamma \partial_{\Theta_\alpha} \partial_{\Theta_\beta} , \quad (\overset{i}{\Delta}_{-1})^2 = 0 ,$$

corresponding to the Lie algebras with structure constants $c_{\alpha\beta}^i{}^\gamma$ ($i = 1, 2$), vanish provided that $c_{\alpha\beta}^i{}^\gamma$ satisfy compatibility conditions [27]

$$\sum_{(\alpha\beta\gamma)} c_{\alpha\beta}^i{}^\lambda c_{\lambda\gamma}^k{}^\delta = 0 ,$$

where $\{ik\}$ denotes the symmetrization of the indices i and k .

The Lie superalgebra (20)-(34), naturally connected with the linear odd Poisson bracket (12), may be useful for the subsequent development of the Batalin-Vilkovisky formalism for the quantization of gauge theories. Indeed, very similar to (12) odd Poisson brackets on the Grassmann algebra are used in a generalization [28] of the triplectic formalism [29] which is a covariant version of the $Sp(2)$ -symmetric quantization [8] of general gauge theories. We should therefore expect that the Lie superalgebra (20)-(34), closely related with the linear odd bracket (12), will also find the application for the further development of the above-mentioned generalization of the triplectic formalism. Let us note that the superalgebra (20)-(34) can also be used in the theory of representations of the semi-simple Lie groups.

The author is sincerely thankful to V.D. Gershun, D.A. Leites and S.L. Lyakhovich for useful discussions and is indebted to J.D. Stasheff for illuminating remarks. The author wishes to thank J. Wess for kind hospitality at the University of Munich where this work was completed.

This work was supported in part by the Ukrainian State Foundation of Fundamental Researches, Grant No 2.5.1/54 and by Grant INTAS No 93-127 (Extension).

References

- [1] V.A. Soroka, Degenerate odd Poisson bracket on Grassmann variables, hep-th/9811223.
- [2] V.A. Soroka, Phys. Lett. B 451, (1999) 349; hep-th/9811252.
- [3] F.A. Berezin, Introduction to algebra and analysis with anticommuting variables, (Moscow State University, 1983).
- [4] M.V. Karasev, V.P. Maslov, Non-linear Poisson brackets. Geometry and quantization, (Moscow, Nauka, 1991).
- [5] J.L. Martin, Proc. Roy. Soc. A 251 (1959) 536.
- [6] I.A. Batalin, G.A. Vilkovisky, Phys. Lett. B 102 (1981) 27.
- [7] I.A. Batalin, G.A. Vilkovisky, Phys. Rev. D 28 (1983) 2567.
- [8] I.A. Batalin, P.M. Lavrov, I.V. Tyutin, J. Math. Phys. 31 (1990) 1487.
- [9] I.A. Batalin, I.V. Tyutin, Int. J. Mod. Phys. A 8 (1993) 2333.
- [10] A.S. Schwarz, Commun. Math. Phys. 155 (1993) 249.
- [11] O.M. Khudaverdian, A.P. Nersessian, Mod. Phys. Lett. A 8 (1993) 2377.

- [12] D.A. Leites, *Docl. Acad. Nauk SSSR* 236 (1977) 804.
- [13] D.V. Volkov, A.I. Pashnev, V.A. Soroka and V.I. Tkach, *JETP Lett.* 44 (1986) 70; *Teor. Mat. Fiz.* 79 (1989) 117.
- [14] V.A. Soroka, *Lett. Math. Phys.* 17 (1989) 201.
- [15] O.M. Khudaverdian, *J. Math. Phys.* 32 (1991) 1934.
- [16] O.M. Khudaverdian, A.P. Nersessian, *J. Math. Phys.* 32 (1991) 1938.
- [17] D.V. Volkov, V.A. Soroka, V.I. Tkach, *Yad. Fiz.* 44 (1986) 810.
- [18] D.V. Volkov, V.A. Soroka, *Yad. Fiz.* 46 (1987) 110.
- [19] V.A. Soroka, *JETP Lett.* 59 (1994) 219.
- [20] V.A. Soroka, Odd Poisson bracket in Hamilton's dynamics, in: *Proceedings of the Workshop on Variational and Local Methods in the Study of Hamiltonian Systems*, (ICTP, Trieste, Italy, 24-28 October, 1994), eds. A. Ambrosetti and G.F. Dell'Antonio (World Scientific, Singapore, 1995) p. 192; hep-th/9503214.
- [21] A.P. Nersessian, *JETP Lett.* 58 (1993) 64.
- [22] D.V. Volkov, A.V. Tur, V.V. Yanovsky, *Phys. Lett. A* 203 (1995) 357.
- [23] V.A. Soroka, *Phys. Atom. Nucl.* 59 (1996) 1270; hep-th/9507030.
- [24] V.A. Soroka, Remark concerning integrable Hamilton systems, in: *Proceedings of the International D. Volkov Memorial Seminar "Supersymmetry and Quantum Field Theory"*, (Kharkov, Ukraine, 5-7 January, 1997), eds. J. Wess, V.P. Akulov. *Lecture Notes in Physics*, Vol. 509, p. 252 (Springer, 1998).
- [25] Z. Popowicz, *Phys. Lett. B* 459, (1999) 150.
- [26] O. Lechtenfeld, A. Sorin, Supersymmetric KP hierarchy in $N = 1$ superspace and its $N = 2$ reductions, solv-int/9907021.
- [27] I.A. Batalin, S.L. Lyakhovich and I.V. Tyutin, *Mod. Phys. Lett. A* 7 (1992) 1931.
- [28] M.A. Grigoriev, *Phys. Lett. B* 458, (1999) 499.
- [29] I.A. Batalin, R. Marnelius and A.M. Semikhatov, *Nucl. Phys. B* 446 (1995) 249.